

National Childhood Immunisation Schedule, Singapore

Vaccination against	Birth	1 Month	3 months	4 months	5 months	6 months	12 months	15 months	18 months	6-7 years [^]	10-11 years ^{^^}
Tuberculosis	BCG										
Hepatitis B	HepB (D1)	HepB (D2)			HepB (D3) [#]						
Diphtheria, Tetanus, Pertussis			DTaP (D1)	DTaP (D2)	DTaP (D3)				DTaP (B1)		Tdap (B2)
Poliovirus			IPV (D1)	IPV (D2)	IPV (D3)				IPV (B1)		OPV (B2)
<i>Haemophilus influenzae</i> type b			Hib (D1)	Hib (D2)	Hib (D3)				Hib (B1)		
Measles, Mumps, Rubella							MMR (D1)	MMR (D2) ^{##}			
Pneumococcal Disease			PCV (D1)		PCV (D2)		PCV (B1)				
Human Papillomavirus		<i>Recommended for females 9 to 26 years; three doses are required at intervals of 0, 2, 6 months</i>									

Notes:

BCG	Bacillus Calmette-Guérin vaccine	#	3 rd dose of HepB can be given at the same time as the 3 rd dose of DTaP, IPV, and Hib for the convenience of parents.
HepB	Hepatitis B vaccine	##	2 nd dose of MMR can be given between 15-18 months
Hib	<i>Haemophilus influenzae</i> type b vaccine		
DTaP	Paediatric diphtheria and tetanus toxoid and acellular pertussis vaccine		
Tdap	Tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine		
MMR	Measles, mumps, and rubella vaccine		
OPV	Oral polio vaccine		
IPV	Inactivated polio vaccine		
PCV	Pneumococcal conjugate vaccine		
D1/D2/D3	1 st dose, 2 nd dose, 3 rd dose		
B1/B2/B3	1 st booster, 2 nd booster, 3 rd booster		
[^]	Primary 1		
^{^^}	Primary 5		